

PRESCRIBING THE SCALAR CURVATURE IN THE NULL CASE

OLIVIER DRUET

ABSTRACT. In this paper, we give a proof of the Kazdan-Warner conjecture concerning the prescribed scalar curvature problem in the null case.

1. INTRODUCTION

Let (M, g) be a smooth compact Riemannian manifold of dimension $n \geq 3$. The Yamabe invariant of (M, g) is a conformal invariant defined by

$$\mu_g = \frac{n-2}{4(n-1)} \inf_{\tilde{g} \in [g]} Vol_{\tilde{g}}(M)^{-\frac{n-2}{n}} \int_M S_{\tilde{g}} dv_{\tilde{g}} \quad (1)$$

where

$$[g] = \{e^u g, u \in C^\infty(M)\}$$

is the conformal class of g , $Vol_{\tilde{g}}(M)$ denotes the volume of M with respect to the Riemannian measure associated to $\tilde{g}$ and $S_{\tilde{g}}$ denotes the scalar curvature of $(M, \tilde{g})$. If $\tilde{g} = u^{\frac{4}{n-2}} g$ for some smooth positive function u , then the scalar curvatures of g and $\tilde{g}$ are related by the following equation :

$$\Delta_g u + \frac{n-2}{4(n-1)} S_g u = \frac{n-2}{4(n-1)} S_{\tilde{g}} u^{2^*-1} \quad (2)$$

where $2^* = \frac{2n}{n-2}$ and $\Delta_g u = -div_g(\nabla u)$. This permits to rewrite (1) as

$$\mu_g = \inf_{u \in C^\infty(M), u \neq 0} \frac{\int_M \left(|\nabla u|_g^2 + \frac{n-2}{4(n-1)} S_g u^2 \right) dv_g}{\left(\int_M |u|^{2^*} dv_g \right)^{\frac{2}{2^*}}}. \quad (3)$$

In this paper, we deal with the so-called prescribed scalar curvature problem in the null case. We assume that $\mu_g = 0$ and we consider the following question : given $f \in C^\infty(M)$, does there exist $\tilde{g} \in [g]$ such that $S_{\tilde{g}} = f$? Thanks to equation (2), and up to some obvious rescaling, the question is equivalent to : given $f \in C^\infty(M)$, does there exist some smooth positive solution u of

$$\Delta_g u + \frac{n-2}{4(n-1)} S_g u = f u^{2^*-1} ?$$

The case $f \equiv 0$ corresponds to the well-known Yamabe problem in the null case, see Yamabe [15], and has been solved in this situation by Trudinger [14]. Thus we may assume without loss of generality that $S_g \equiv 0$ and that $Vol_g(M) = 1$ since there always exists such a metric in the conformal class of g . Then the above equation becomes

$$\Delta_g u = f u^{2^*-1}. \quad (4)$$

Date: July, 2005.

Obvious necessary conditions on f so that there exists a smooth positive solution u of the above equation are that

$$\int_M f dv_g < 0 \quad (5)$$

and that

$$\max_M f > 0 \quad (6)$$

if $f \not\equiv 0$, what we will assume in the sequel since the case $f \equiv 0$ is already settled. In order to prove that (5) is a necessary condition, one has just to multiply equation (4) by u^{1-2^*} and to integrate over M to obtain that

$$\int_M f dv_g = \int_M u^{1-2^*} \Delta_g u dv_g = -\frac{n+2}{n-2} \int_M u^{-2^*} |\nabla u|_g^2 dv_g < 0$$

since $u \not\equiv 0$ if $f \not\equiv 0$. As for the second necessary condition (6), one has just to integrate equation (4) over M to obtain that

$$\int_M f u^{2^*-1} dv_g = 0$$

which enforces f to change sign if it is not identically zero.

The natural question then is : are the conditions (5) and (6) sufficient for f to be the scalar curvature of a metric conformal to g ? In the sequel of their 2-dimensional work [12], Kazdan and Warner conjectured that the answer to this question was positive (see [11, 13]). Jung, in [10], used the technique of sub- and upper-solution to solve equation (4) under the assumptions (5) and (6). However, there is some problem in this proof - see p. 743 of [10] - since the limit, for $s > 0$ real, of

$$\left(\frac{1 - x^x}{s^{x^2} - x^x} \right)^{-\frac{\ln x}{x}}$$

as $x \rightarrow 0^+$ is not 1 but s^{-1} . We provide in this paper a proof of the conjecture of Kazdan and Warner, based on fine blow-up analysis. In other words, we prove the following result :

Theorem 1.1. *Let (M, g) be a smooth compact Riemannian manifold of dimension $n \geq 3$ with null scalar curvature. A smooth function f on M is the scalar curvature of a metric conformal to g if and only if $f \equiv 0$ or (5) and (6) hold.*

Note that this result had already been proved in dimensions 3 and 4 by minimisation techniques and test functions computations by Escobar and Schoen [6]. In this work, they also proved the existence of a solution of the equation (4) under some flatness assumptions on f around one of its maximum point in the case (M, g) is locally conformally flat.

The solution we obtain during the proof of the theorem is a solution of minimal energy, a so-called minimizing solution.

Let us at last remark that, given (M, g) a smooth compact Riemannian manifold of dimension $n \geq 3$ with null Yamabe invariant, there exists a unique $\tilde{g} = \varphi^{\frac{4}{n-2}} g$ with zero scalar curvature and $Vol_{\tilde{g}}(M) = 1$. The conditions (5) and (6) of theorem 1.1 can be rewritten in terms of g as follows : $\max_M f > 0$ and $\int_M f \varphi^{2^*} dv_g < 0$.

2. PROOF OF THE THEOREM

We assume in the following that (M, g) has null scalar curvature and that $Vol_g(M) = 1$. We let $f \in C^\infty(M)$, $f \neq 0$, be such that (5) and (6) hold. We first recall, and give a quick proof of, the following classical result which deals with the subcritical equation :

Claim 2.1. *Let (M, g) be a smooth compact Riemannian manifold of dimension $n \geq 3$ with null scalar curvature. Let $f \in C^\infty(M)$, $f \neq 0$, be such that (5) and (6) hold. For any $2 < q < 2^*$, there exists $u_q \in C^\infty(M)$, $u_q > 0$, such that*

$$\int_M |\nabla u_q|_g^2 dv_g = \lambda_q, \quad u_q \in \mathcal{H}_q$$

where

$$\mathcal{H}_q = \left\{ u \in H_1^2(M) \text{ s.t. } \int_M f|u|^q dv_g = 1 \right\}$$

and

$$\lambda_q = \inf_{u \in \mathcal{H}_q} \int_M |\nabla u|_g^2 dv_g.$$

Moreover, u_q satisfies that

$$\Delta_g u_q = \lambda_q f u_q^{q-1}.$$

Proof - It is rather standard and we just outline the proof. First, under the assumptions made on f , it is easily checked that $\mathcal{H}_q$ is not empty. Let (u_i) be a sequence of functions in $\mathcal{H}_q$ such that

$$\int_M |\nabla u_i|_g^2 dv_g \rightarrow \lambda_q$$

as $i \rightarrow +\infty$. If we prove that (u_i) is bounded in $H_1^2(M)$, then we can extract a subsequence which converges to a solution of our problem. It is straightforward since the embedding of $H_1^2(M)$ into $L^q(M)$ is compact for $q < 2^*$. Moreover, our minimizer may be choosen nonnegative and the result will follow from the Euler-Lagrange equation satisfied by the minimizer, from standard regularity theory and from the maximum principle. Thus it remains to prove that (u_i) is bounded in $H_1^2(M)$. Since $(|\nabla u_i|_g)$ is bounded in $L^2(M)$, we just have to prove that (u_i) is bounded in $L^2(M)$. Assume by contradiction that $\|u_i\|_2 \rightarrow +\infty$ as $i \rightarrow +\infty$. Let $v_i = \frac{u_i}{\|u_i\|_2}$. Then it is clear that (v_i) is bounded in $H_1^2(M)$ and that $\|\nabla v_i\|_2 \rightarrow 0$ as $i \rightarrow +\infty$. Thus $v_i \rightarrow v_\infty$ strongly in $H_1^2(M)$ as $i \rightarrow +\infty$ where $v_\infty \equiv 1$. Remember that $Vol_g(M) = 1$. We thus obtain that

$$0 = \lim_{i \rightarrow +\infty} \int_M f |v_i|^q dv_g = \int_M f |v_\infty|^q dv_g = \int_M f dv_g,$$

which is a contradiction since $\int_M f dv_g < 0$. This ends the proof of the claim. $\diamond$

Let us now consider the solution u_q of

$$\Delta_g u_q = \lambda_q f u_q^{q-1}, \quad \int_M f u_q^q dv_g = 1, \quad (7)$$

given by claim 2.1. Note that $\lambda_q > 0$ since, otherwise, u_q would be constant and the constraint $\int_M f u_q^q dv_g = 1$ could not be satisfied since $\int_M f dv_g < 0$. If (u_q)

is bounded in $L^\infty(M)$, then standard elliptic theory gives that, after passing to a subsequence, $u_q \rightarrow u_0$ in $C^2(M)$ as $q \rightarrow 2^*$ where u_0 satisfies that

$$\Delta_g u_0 = \lambda_{2^*} f u_0^{2^*-1}, \quad \int_M f u_0^{2^*} dv_g = 1.$$

Indeed, it is easy to check that

$$\lim_{q \rightarrow 2^*} \lambda_q = \lambda_{2^*}. \quad (8)$$

Then it is clear that $\lambda_{2^*} > 0$ and that $\lambda_{2^*}^{\frac{4}{n-2}} u_0$ is a smooth positive solution of (4) thanks to the maximum principle and standard regularity theory. Thus, in this case, the theorem is proved. We assume now by contradiction that

$$u_q(x_q) = \|u_q\|_\infty \rightarrow +\infty \text{ as } q \rightarrow 2^*. \quad (9)$$

We prove the following :

Claim 2.2. *Assume that (9) holds. Then we have that, after passing to a subsequence,*

$$u_q(x_q)^{-1} u_q \left(\exp_{x_q}(\mu_q x) \right) \rightarrow (1 + |x|^2)^{1-\frac{n}{2}}$$

in $C_{loc}^2(\mathbb{R}^n)$ as $q \rightarrow 2^*$ where

$$\mu_q = \left(\frac{n(n-2)}{\lambda_q f(x_q)} \right)^{\frac{1}{2}} u_q(x_q)^{-\frac{q-2}{2}}.$$

Proof - Let us first remark that (u_q) is bounded in $H_1^2(M)$. Since $\|\nabla u_q\|_2^2 \rightarrow \lambda_{2^*}$ as $q \rightarrow 2^*$, we only need to prove that (u_q) is bounded in $L^2(M)$. Assume on the contrary that $\|u_q\|_2 \rightarrow +\infty$ as $q \rightarrow 2^*$. We then easily obtain that

$$\frac{u_q}{\|u_q\|_2} \rightarrow 1$$

strongly in $H_1^2(M)$ as $q \rightarrow 2^*$. This is clearly in contradiction with the fact that $\int_M f u_q^q dv_g = 1$ since $\int_M f dv_g < 0$. Thus (u_q) is bounded in $H_1^2(M)$. As a consequence of Sobolev's embeddings, we know that there exists some $\Lambda > 0$ such that

$$\int_M u_q^q dv_g \leq \Lambda. \quad (10)$$

We let now $\delta > 0$ small enough and we set for $x \in B_0 \left(\delta u_q(x_q)^{\frac{q-2}{2}} \right)$

$$\begin{aligned} v_q(x) &= u_q(x_q)^{-1} u_q \left(\exp_{x_q} \left(u_q(x_q)^{-\frac{q-2}{2}} x \right) \right), \\ g_q(x) &= \exp_{x_q}^* g \left(\exp_{x_q} \left(u_q(x_q)^{-\frac{q-2}{2}} x \right) \right), \\ f_q(x) &= f \left(\exp_{x_q} \left(u_q(x_q)^{-\frac{q-2}{2}} x \right) \right). \end{aligned} \quad (11)$$

It is clear that $g_q \rightarrow \xi$, the Euclidean metric, in $C_{loc}^2(\mathbb{R}^n)$ as $q \rightarrow 2^*$, that $f_q \rightarrow f(x_0)$ in $C_{loc}^2(\mathbb{R}^n)$ as $q \rightarrow 2^*$ where, after passing to a subsequence, $x_0 = \lim_{q \rightarrow 2^*} x_q$ and that $0 \leq v_q \leq 1$ in $B_0 \left(\delta u_q(x_q)^{\frac{q-2}{2}} \right)$ with $v_q(0) = 1$. Moreover, we have that

$$\Delta_{g_q} v_q = \lambda_q f_q v_q^{q-1}$$

in $B_0 \left(\delta u_q(x_q)^{\frac{q-2}{2}} \right)$. Standard elliptic theory gives then that

$$v_q \rightarrow v_0 \text{ in } C_{loc}^2(\mathbb{R}^n) \text{ as } q \rightarrow 2^*$$

where $0 \leq v_0 \leq 1$, $v_0(0) = 1$ and

$$\Delta_\xi v_0 = \lambda_{2^*} f(x_0) v_0^{2^*-1}.$$

Since 0 is a point of maximum of v_0 , $\Delta_\xi v_0(0) \geq 0$ and thus $f(x_0) \geq 0$. Assume by contradiction that

$$\lambda_{2^*} f(x_0) = 0. \quad (12)$$

Then v_0 is a bounded harmonic function in $\mathbb{R}^n$ with $v_0(0) = 1$ and thus $v_0 \equiv 1$. Now we write thanks to (10) that

$$\begin{aligned} \Lambda &\geq \int_{B_{x_q}} \left(R u_q(x_q)^{-\frac{q-2}{2}} \right) u_q^q dv_g \\ &\geq u_q(x_q)^{(2^*-q)\frac{n-2}{2}} \int_{B_0(R)} v_q^q dv_{g_q} \\ &\geq \int_{B_0(R)} v_0^{2^*} dv_\xi + o(1). \end{aligned}$$

This clearly proves that $v_0 \in L^{2^*}(\mathbb{R}^n)$ and thus that $v_0 \not\equiv 1$. In particular, (12) can not hold and $\lambda_{2^*} f(x_0) > 0$. It is then clear that

$$u_q(x_q)^{-1} u_q \left(\exp_{x_q}(\mu_q x) \right) \rightarrow U_0$$

in $C_{loc}^2(\mathbb{R}^n)$ as $q \rightarrow 2^*$ where μ_q is as in the proposition and U_0 satisfies that $0 \leq U_0 \leq 1$, $U_0(0) = 1$ and

$$\Delta_\xi U_0 = n(n-2)U_0^{2^*-1}.$$

By the classification result of Caffarelli-Gidas-Spruck [3], we know that

$$U_0(x) = (1 + |x|^2)^{1-\frac{n}{2}}.$$

This ends the proof of the claim. $\diamond$

We next prove the following :

Claim 2.3. *Under the assumption (9), we have that, after passing to a subsequence, $u_q \rightharpoonup 0$ in $H_1^2(M)$ as $q \rightarrow 2^*$ and that*

$$\lim_{q \rightarrow 2^*} \lambda_q = \lambda_{2^*} = \left(\max_M f \right)^{-\frac{2}{2^*}} K(n, 2)^{-2}$$

where

$$K(n, 2)^2 = \frac{4}{n(n-2)} \omega_n^{-\frac{2}{n}}$$

with ω_n the volume of the unit n -sphere in $\mathbb{R}^{n+1}$. Moreover, we have that

$$\lim_{R \rightarrow +\infty} \lim_{q \rightarrow 2^*} \int_{B_{x_q}(R\mu_q)} f u_q^q dv_g = 1,$$

that $f(x_0) = \max_M f$ and that

$$\lim_{q \rightarrow 2^*} u_q(x_q)^{2^*-q} = 1.$$

Proof - Thanks to claim 2.2, we can write that

$$\begin{aligned} & \int_{B_{x_q}(R\mu_q)} f u_q^q dv_g \\ &= f(x_0)^{1-\frac{n}{2}} (n(n-2))^{\frac{n}{2}} \lambda_q^{-\frac{n}{2}} u_q(x_q)^{(2^*-q)\frac{n-2}{2}} \left(\int_{B_0(R)} U_0^{2^*} dx + o(1) \right). \end{aligned}$$

As shown during the proof of claim 2.2, we have that (u_q) is bounded in $H_1^2(M)$. Thus, by Sobolev's embeddings, (u_q) is uniformly bounded in $L^q(M)$. We deduce that, after passing to a subsequence,

$$u_q(x_q)^{(2^*-q)\frac{n-2}{2}} \rightarrow A \quad (13)$$

as $q \rightarrow 2^*$ for some $A \geq 1$. It is also easily checked that

$$\lim_{q \rightarrow 2^*} \lambda_q = \lambda_{2^*}$$

and that

$$\int_{\mathbb{R}^n} U_0^{2^*} dx = (n(n-2))^{-\frac{n}{2}} K(n, 2)^{-n}.$$

This leads to

$$\lim_{R \rightarrow +\infty} \lim_{q \rightarrow 2^*} \int_{B_{x_q}(R\mu_q)} f u_q^q dv_g = f(x_0)^{1-\frac{n}{2}} \lambda_{2^*}^{-\frac{n}{2}} K(n, 2)^{-n} A. \quad (14)$$

Since (u_q) is bounded in $H_1^2(M)$, after passing to a subsequence, $u_q \rightharpoonup u_0$ weakly in $H_1^2(M)$ as $q \rightarrow 2^*$. It is clear thanks to (14) that the convergence can not be strong. We then write that

$$\int_M f |u_q - u_0|^q dv_g = 1 - \int_M f |u_0|^q dv_g + o(1) \quad (15)$$

and that

$$\int_M |\nabla(u_q - u_0)|_g^2 = \lambda_{2^*} - \int_M |\nabla u_0|_g^2 dv_g + o(1). \quad (16)$$

Equation (16) is trivial while equation (15) is easy to obtain (see for instance [2]). By the definition of λ_q , we have that

$$\lambda_q \left| \int_M f |u_0|^q dv_g \right|^{\frac{2}{q}-1} \int_M f |u_0|^q dv_g \leq \|\nabla u_0\|_2^2$$

and that

$$\lambda_q \left| \int_M f |u_q - u_0|^q dv_g \right|^{\frac{2}{q}-1} \int_M f |u_q - u_0|^q dv_g \leq \|\nabla(u_q - u_0)\|_2^2.$$

Passing to the limit in these two inequalities, using (15) and (16), we get that

$$\lambda_{2^*} |X|^{\frac{2}{2^*}-1} X \leq \|\nabla u_0\|_2^2 \quad (17)$$

and that

$$\lambda_{2^*} |1 - X|^{\frac{2}{2^*}-1} (1 - X) \leq \lambda_{2^*} - \|\nabla u_0\|_2^2 \quad (18)$$

where

$$X = \int_M f |u_0|^{2^*} dv_g.$$

Since $\|\nabla u_0\|_2^2 \leq \lambda_{2^*}$ thanks to (16), the first inequality ensures that $X \leq 1$ while the second inequality ensures that $X \geq 0$. Next we sum (17) and (18) to get that

$$|X|^{\frac{2}{2^*}-1} X + |1-X|^{\frac{2}{2^*}-1} (1-X) \leq 1.$$

Since $0 \leq X \leq 1$, this is possible if and only if $X = 0$ or $X = 1$. But, if $X = 1$, (16) and (17) imply that $u_q \rightarrow u_0$ strongly in $H_1^2(M)$. As already said, this can not happen. Thus $X = 0$ and (18) implies in turn that $\|\nabla u_0\|_2 = 0$ so that $u_0 \equiv Cst$. Since $X = \int_M f |u_0|^{2^*} dx = 0$ and $\int_M f dv_g < 0$, this clearly implies that $u_0 \equiv 0$. This proves the first part of the claim. Thanks to the work of Hebey-Vaugon [8, 9], we know that there exists $B > 0$ such that

$$\|u_q\|_{2^*}^2 \leq K(n, 2)^2 \|\nabla u_q\|_2^2 + B \|u_q\|_2^2$$

for all $q < 2^*$ where $K(n, 2)$ is as in the statement of the claim. Since the embedding of $H_1^2(M)$ into $L^2(M)$ is compact, we have that $\|u_q\|_2 \rightarrow 0$ as $q \rightarrow 2^*$ so that we get that

$$\|u_q\|_{2^*}^2 \leq K(n, 2)^2 \|\nabla u_q\|_2^2 + o(1).$$

In particular, we deduce that

$$\begin{aligned} 1 + \int_M f^- u_q^q dv_g &= \int_M f^+ u_q^q dv_g \\ &\leq \left(\max_M f \right) \|u_q\|_{2^*}^q \text{Vol}_g(M)^{\frac{(2^*-q)}{2^*}} \\ &\leq \left(\max_M f \right) K(n, 2)^q \|\nabla u_q\|_2^q + o(1) \\ &\leq \left(\max_M f \right)^{\frac{2}{2^*}} K(n, 2)^{2^*} \lambda_{2^*}^{\frac{2^*}{2}} + o(1) \end{aligned}$$

where $f^+ = \frac{1}{2}(f + |f|)$ and $f^- = \frac{1}{2}(|f| - f)$. This leads to

$$\lambda_{2^*} \geq \left(\max_M f \right)^{-\frac{2}{2^*}} K(n, 2)^{-2}.$$

Standard test-functions computations, we refer to [6] for instance, show that the reverse inequality also holds. Thus we have that

$$\lambda_{2^*} = \left(\max_M f \right)^{-\frac{2}{2^*}} K(n, 2)^{-2}. \quad (19)$$

Then we can also deduce that

$$\int_M f^- u_q^q dv_g \rightarrow 0 \text{ as } q \rightarrow 2^*. \quad (20)$$

Let us come back to (14) with (19). We obtain that

$$\lim_{R \rightarrow +\infty} \lim_{q \rightarrow 2^*} \int_{B_{x_q}(R\mu_q)} f u_q^q dv_g = \left(\frac{\max_M f}{f(x_0)} \right)^{\frac{n-2}{2}} A.$$

Since $\int_M f u_q^q dv_g = 1$ and thanks to (20), we deduce that $f(x_0) = \max_M f$ and that $A = 1$. This ends the proof of the claim. $\diamond$

We now prove the following weak pointwise estimates on u_q :

Claim 2.4. *There exists $C > 0$ such that*

$$d_g(x_q, x)^{\frac{2}{q-2}} u_q(x) \leq C$$

for all $x \in M$ and all $q < 2^$. Moreover, we have that*

$$\lim_{R \rightarrow +\infty} \lim_{q \rightarrow 2^*} \sup_{M \setminus B_{x_q}(R\mu_q)} d_g(x_q, x)^{\frac{2}{q-2}} u_q(x) = 0.$$

Proof - The proof is rather standard and follows the lines of [4]. We also refer to [5] for this kind of argument. We briefly sketch the proof in the following. Let us set

$$w_q(x) = d_g(x_q, x)^{\frac{2}{q-2}} u_q(x)$$

and let $y_q \in M$ be such that

$$w_q(y_q) = \max_M w_q.$$

Assume by contradiction that

$$w_q(y_q) \rightarrow +\infty \text{ as } q \rightarrow 2^*. \quad (21)$$

Since M is compact, it is clear that $u_q(y_q) \rightarrow +\infty$ as $q \rightarrow 2^*$. We let now $\delta > 0$ small enough and we set for $x \in B_0\left(\delta u_q(y_q)^{\frac{q-2}{2}}\right)$

$$\begin{aligned} v_q(x) &= u_q(y_q)^{-1} u_q\left(\exp_{y_q}\left(u_q(y_q)^{-\frac{q-2}{2}} x\right)\right), \\ g_q(x) &= \exp_{y_q}^* g\left(\exp_{y_q}\left(u_q(y_q)^{-\frac{q-2}{2}} x\right)\right), \\ f_q(x) &= f\left(\exp_{y_q}\left(u_q(y_q)^{-\frac{q-2}{2}} x\right)\right). \end{aligned} \quad (22)$$

It is clear that $g_q \rightarrow \xi$ and that $f_q \rightarrow f(y_0)$ in $C_{loc}^2(\mathbb{R}^n)$ as $q \rightarrow 2^*$ where, after passing to a subsequence, $y_0 = \lim_{q \rightarrow 2^*} y_q$. Moreover, we have that

$$\Delta_{g_q} v_q = \lambda_q f_q v_q^{q-1}$$

in $B_0\left(\delta u_q(y_q)^{\frac{q-2}{2}}\right)$. Let now $x \in \mathbb{R}^n$ and let us write thanks to the choice of y_q we made that

$$d_g\left(x_q, \exp_{y_q}\left(u_q(y_q)^{-\frac{q-2}{2}} x\right)\right) v_q(x)^{\frac{q-2}{2}} \leq d_g(x_q, y_q).$$

This leads to

$$\left(d_g(x_q, y_q) - |x| u_q(y_q)^{-\frac{q-2}{2}}\right) v_q(x)^{\frac{q-2}{2}} \leq d_g(x_q, y_q)$$

which can be written as

$$\left(1 - |x| w_q(y_q)^{-\frac{q-2}{2}}\right) v_q(x)^{\frac{q-2}{2}} \leq 1.$$

We deduce from (21) that

$$v_q(x) \leq 1 + o(1).$$

Standard elliptic theory gives then that

$$v_q \rightarrow v_0 \text{ in } C_{loc}^2(\mathbb{R}^n) \text{ as } q \rightarrow 2^*$$

where $0 \leq v_0 \leq 1$, $v_0(0) = 1$ and

$$\Delta_\xi v_0 = \lambda_{2^*} f(y_0) v_0^{2^*-1}.$$

Mimicking the proof of claim 2.2, one proves that $f(y_0) > 0$. We then write since $\|u_q\|_\infty^{2^*-q} \rightarrow 1$ as $q \rightarrow 2^*$, see claim 2.3, that

$$\lim_{q \rightarrow 2^*} \int_{B_{y_q}} \left(R u_q(y_q)^{-\frac{q-2}{2}} \right) f u_q^q dv_g = f(y_0) \int_{B_0(R)} v_0^{2^*} dx > 0. \quad (23)$$

Note that (21) implies that

$$\frac{d_g(x_q, y_q)}{u_q(y_q)^{-\frac{q-2}{2}}} \rightarrow +\infty$$

and that

$$\frac{d_g(x_q, y_q)}{\mu_q} \rightarrow +\infty$$

as $q \rightarrow 2^*$ thanks to the definition of μ_q and of x_q . Thus we can write thanks to (20) that for any $R > 0$,

$$\int_M f u_q^q dv_g \geq \int_{B_{x_q}(R\mu_q)} f u_q^q dv_g + \int_{B_{y_q}} \left(R u_q(y_q)^{-\frac{q-2}{2}} \right) f u_q^q dv_g + o(1)$$

for q close to 2^* which leads to a contradiction thanks to claim 2.3 and to (23) since $\int_M f u_q^q dv_g = 1$. Thus we have proved the first part of the claim. Note that, as a consequence of this estimate, we know that (u_q) is uniformly bounded in any compact subset of $M \setminus \{x_0\}$. Standard elliptic theory then gives since $u_q \rightarrow 0$ in $H_1^2(M)$ that

$$u_q \rightarrow 0 \text{ in } C_{loc}^2(M \setminus \{x_0\}) \text{ as } q \rightarrow 2^*. \quad (24)$$

Let us now assume by contradiction that the second estimate of the proposition does not hold. In other words, let us assume that there exists $z_q \in M$ such that

$$\frac{d_g(x_q, z_q)}{\mu_q} \rightarrow +\infty \text{ as } q \rightarrow 2^* \quad (25)$$

and that

$$w_q(z_q) \geq \varepsilon_0 \quad (26)$$

for some $\varepsilon_0 > 0$. Note that (26) together with (24) implies that $d_g(x_q, z_q) \rightarrow 0$ as $q \rightarrow 2^*$ and thus that $u_q(z_q) \rightarrow +\infty$ as $q \rightarrow 2^*$. One can then check that, after passing to a subsequence,

$$u_q(z_q)^{-1} u_q \left(\exp_{z_q} \left(u_q(z_q)^{-\frac{q-2}{2}} x \right) \right) \rightarrow V_0$$

in $C_{loc}^2(\mathbb{R}^n \setminus \{P\})$ where

$$P = \lim_{q \rightarrow 2^*} u_q(z_q)^{\frac{q-2}{2}} \exp_{z_q}^{-1}(x_q)$$

which does exist and satisfies $\varepsilon_0^{\frac{2}{2^*-2}} \leq |P| \leq C^{\frac{2}{2^*-2}}$ where C is the constant involved in the first estimate of the claim. Moreover, V_0 satisfies that $V_0(0) = 1$ and that

$$\Delta_\xi V_0 = \lambda_{2^*} f(x_0) V_0^{2^*-1} \text{ in } \mathbb{R}^n \setminus \{P\}.$$

This implies in particular that

$$\lim_{q \rightarrow 2^*} \int_{B_{z_q}} \left(\frac{1}{\frac{1}{2}\varepsilon_0^{\frac{2}{2^*-2}}} u_q(z_q)^{-\frac{q-2}{2}} \right) f u_q^q dv_g = f(x_0) \int_{B_0} \left(\frac{1}{\frac{1}{2}\varepsilon_0^{\frac{2}{2^*-2}}} \right) V_0^{2^*} dx > 0.$$

Using (25) and (26), one proves that, for any $R > 0$, the balls $B_{x_q}(R\mu_q)$ and $B_{z_q}\left(\frac{1}{2}\varepsilon_0^{\frac{2}{2^*-2}}u_q(z_q)^{-\frac{q-2}{2}}\right)$ are disjoint for q close to 2^* . And one obtains a contradiction as in the first part of the proof. This ends the proof of the claim. $\diamond$

We transform now this weak pointwise estimate into an almost optimal pointwise estimate. Once again, we refer to [5] for the general scheme of this kind of proof.

Claim 2.5. *For any $0 < \varepsilon < 1$, there exists $C_\varepsilon > 0$ such that*

$$u_q(x) \leq C_\varepsilon \left(\mu_q^{\frac{n-2}{2}(1-2\varepsilon)} d_g(x_q, x)^{(2-n)(1-\varepsilon)} + \eta_q d_g(x_q, x)^{(2-n)\varepsilon} \right)$$

for all $x \in M \setminus \{x_q\}$ and all $q < 2^*$. Here, η_q is defined by

$$\eta_q = \sup_{M \setminus B_{x_q}(\delta)} u_q$$

where $\delta > 0$ is fixed small enough.

Proof - We let G be the Green function for the Laplacian on M normalized such that

$$\int_M G(x, y) dv_g(y) = 0$$

for all $x \in M$. We let $G_q(x) = G(x_q, x)$ and we let $\delta > 0$ be such that

$$d_g(x_q, x) \frac{|\nabla G_q(x)|}{G_q(x)} \geq \frac{n-2}{2} \quad \text{and} \quad G_q(x) \geq 1 \quad (27)$$

for $x \in B_x(2\delta)$. For the construction of and estimates on the Green function, we refer the reader to the Appendix of [5]. We fix $0 < \varepsilon < 1$ and we choose $R_\varepsilon > 0$ such that

$$\lambda_q d_g(x_q, x)^2 u_q(x)^{q-2} \leq \frac{\varepsilon(1-\varepsilon)}{f(x_0)} \frac{(n-2)^2}{8} \quad \text{in } M \setminus B_{x_q}(R_\varepsilon \mu_q) \quad (28)$$

for q close to 2^* thanks to claim 2.4. We let L_q be the linear operator defined by

$$L_q \varphi = \Delta_g \varphi - \lambda_q f u_q^{q-2} \varphi.$$

This operator satisfies the maximum principle on M in the following sense (see [1]) : if Ω is a smooth subset of M and if φ and ψ are two smooth functions in a neighbourhood of Ω such that

$$L_q \varphi \geq L_q \psi \quad \text{in } \Omega$$

and

$$\varphi \geq \psi \quad \text{on } \partial\Omega,$$

then $\varphi \geq \psi$ in Ω . We note that $L_q u_q = 0$ in M . Simple computations lead thanks to (27) and (28) to

$$\begin{aligned} \frac{L_q G_q^\nu}{G_q^\nu} &= \nu(1-\nu) \frac{|\nabla G_q|^2}{G_q^2} - \lambda_q f u_q^{q-2} \\ &\geq d_g(x_q, x)^{-2} \left(\nu(1-\nu) \frac{(n-2)^2}{4} - \lambda_q f(x_0) d_g(x_q, x)^2 u_q(x)^{q-2} \right) \\ &\geq 0 \end{aligned}$$

in $B_{x_q}(2\delta) \setminus B_{x_q}(R_\varepsilon \mu_q)$ for q close to 2^* and for $\nu = \varepsilon$ and $\nu = 1 - \varepsilon$. Using now claim 2.2 and standard estimates on the Green function, we obtain the existence of some $C_\varepsilon > 0$ such that

$$u_q \leq C_\varepsilon \left(\mu_q^{\frac{n-2}{2}(1-2\varepsilon)} G_q(x)^{1-\varepsilon} + \eta_q G_q(x)^\varepsilon \right)$$

on $\partial(B_{x_q}(2\delta) \setminus B_{x_q}(R_\varepsilon \mu_q))$. Note that we used the fact that $u_q(x_q)^{2^*-q} \rightarrow 1$ as $q \rightarrow 2^*$, which was proved in claim 2.3. The maximum principle and standard estimates on the Green function permit to conclude the proof of the claim. Note that, outside $B_{x_q}(2\delta)$, the estimate of the claim is obviously satisfied, up to change C_ε . $\diamond$

Remark that, by standard elliptic theory, see for instance [7], chapter 8, we know that

$$\frac{u_q}{\eta_q} \rightarrow H$$

in $C_{loc}^2(M \setminus \{x_0\})$ as $q \rightarrow 2^*$ where H is a nonzero positive function satisfying that $\Delta_g H = 0$ in $M \setminus \{x_0\}$. Multiplying the equation (7) by u_q^{1-q} and integrating over M , we obtain that

$$\begin{aligned} \lambda_q \int_M f dv_g &= -(q-1) \int_M u_q^{-q} |\nabla u_q|_g^2 dv_g \\ &\leq -(q-1) \eta_q^{2-q} \left(\int_{M \setminus B_{x_0}(\delta)} H^{-q} |\nabla H|^2 dv_g + o(1) \right) \end{aligned}$$

for all $\delta > 0$. Since $\eta_q \rightarrow 0$ as $q \rightarrow 2^*$ thanks to (24), this clearly implies that $\nabla H \equiv 0$. Thus, by the definition of η_q , we get that $H \equiv 1$ and we have proved that

$$\frac{u_q}{\eta_q} \rightarrow 1 \text{ in } C_{loc}^2(M \setminus \{x_0\}) \text{ as } q \rightarrow 2^*. \quad (29)$$

We now describe precisely the asymptotic behaviour of u_q :

Claim 2.6. *For any sequence (y_q) of points in M , we have that*

$$\begin{aligned} u_q(y_q) &= \overline{u_q} + 2^n \frac{\omega_{n-1}}{n} u_q(x_q)^{-1} \varphi(y_q) + o(u_q(x_q)^{-1}) \\ &\quad + u_q(x_q) \left(1 + \frac{d_g(x_q, y_q)^2}{\mu_q^2} \right)^{-\frac{n-2}{2}} (\Phi(x_q, y_q) + o(1)). \end{aligned}$$

In this equation,

$$\overline{u_q} = \int_M u_q dv_g$$

and φ is a solution of

$$\Delta_g \varphi = \lambda_{2^*} \left(1 - \frac{f}{\bar{f}} \right)$$

with $\bar{\varphi} = 0$ and

$$\Phi(x, y) = (n-2) \omega_{n-1} d_g(x, y)^{n-2} G(x, y)$$

where G is the Green function of the Laplacian on M normalized such that

$$\int_M G(x, y) dv_g(y) = 0$$

for all $x \in M$.

Proof - We fix $0 < \varepsilon < \frac{2}{n+2}$. We first write by integrating equation (7) over M that

$$\int_M f u_q^{q-1} dv_g = 0$$

and thus that

$$\int_{\{u_q(x) > \alpha \eta_q\}} f u_q^{q-1} dv_g = - \int_{\{u_q(x) \leq \alpha \eta_q\}} f u_q^{q-1} dv_g \quad (30)$$

for all $\alpha > 1$. Thanks to Lebesgue's dominated convergence theorem and to (29), we have that

$$\int_{\{u_q(x) \leq \alpha \eta_q\}} f u_q^{q-1} dv_g = \eta_q^{q-1} \int_M f dv_g + o(\eta_q^{q-1}) . \quad (31)$$

We write also thanks to claim 2.2 that, for any $R > 0$,

$$\begin{aligned} \int_{\{u_q(x) > \alpha \eta_q\}} f u_q^{q-1} dv_g &= \int_{B_{x_q}(R\mu_q)} f u_q^{q-1} dv_g \\ &\quad + \int_{\{u_q(x) > \alpha \eta_q\} \setminus B_{x_q}(R\mu_q)} f u_q^{q-1} dv_g . \end{aligned} \quad (32)$$

Thanks to claim 2.5, we have that

$$\begin{aligned} &\left| \int_{\{u_q(x) > \alpha \eta_q\} \setminus B_{x_q}(R\mu_q)} f u_q^{q-1} dv_g \right| \\ &\leq C \mu_q^{\frac{n-2}{2}(1-2\varepsilon)(q-1)} \int_{M \setminus B_{x_q}(R\mu_q)} d_g(x_q, x)^{(2-n)(1-\varepsilon)(q-1)} dv_g \\ &\quad + C \eta_q^{q-1} \int_{\{u_q(x) > \alpha \eta_q\} \setminus B_{x_q}(R\mu_q)} d_g(x_q, x)^{(2-n)\varepsilon(q-1)} dv_g \\ &\leq C R^{n-(n+2)(1-\varepsilon)} \mu_q^{\frac{n-2}{2}} + C \alpha^{-\frac{n-(n+2)\varepsilon}{(n-2)\varepsilon}} \eta_q^{q-1} + C \alpha^{\frac{1-2\varepsilon}{\varepsilon}(q-1)} \mu_q^{\frac{n+2}{2}(1-2\varepsilon)} \end{aligned}$$

where C changes from line to line but is always independent of R , α and q . Here we used the fact that $\varepsilon < \frac{2}{n+2}$ and the fact that $u_q(x_q)^{2^*-q} \rightarrow 1$ as $q \rightarrow 2^*$, which was proved in claim 2.3. We also used the following consequence of claim 2.5 :

$$\begin{aligned} \text{Either } u_q(x) > \alpha \eta_q &\Rightarrow d_g(x_q, x)^{(n-2)\varepsilon} \leq \frac{2C_\varepsilon}{\alpha} , \\ \text{or } \eta_q &\leq \left(\frac{\alpha}{2C_\varepsilon} \right)^{\frac{1-2\varepsilon}{\varepsilon}} \mu_q^{\frac{n-2}{2}(1-2\varepsilon)} . \end{aligned} \quad (33)$$

Since $0 < \varepsilon < \frac{2}{n+2}$, we deduce from the above estimate and the definition of μ_q that

$$\lim_{\alpha \rightarrow +\infty} \lim_{R \rightarrow +\infty} \lim_{q \rightarrow 2^*} \frac{\int_{\{u_q(x) > \alpha \eta_q\} \setminus B_{x_q}(R\mu_q)} f u_q^{q-1} dv_g}{u_q(x_q)^{-1} + \eta_q^{q-1}} = 0 . \quad (34)$$

Now, thanks to claims 2.2 and 2.3, it is easily checked that

$$\lim_{R \rightarrow +\infty} \lim_{q \rightarrow 2^*} u_q(x_q) \int_{B_{x_q}(R\mu_q)} f u_q^{q-1} dv_g = \frac{2^n \omega_{n-1}}{n} . \quad (35)$$

Coming back to (32) with (34) and (35), we thus get that

$$\int_{\{u_q(x) > \alpha \eta_q\}} f u_q^{q-1} dv_g = \frac{2^n \omega_{n-1}}{n} u_q(x_q)^{-1} + o(u_q(x_q)^{-1}) + o(\eta_q^{q-1}) . \quad (36)$$

Coming back to (30) with (31) and (36), we obtain that

$$\eta_q^{q-1} = -\frac{2^n \omega_{n-1}}{n \bar{f}} u_q(x_q)^{-1} + o\left(u_q(x_q)^{-1}\right). \quad (37)$$

We write now thanks to the Green representation formula that

$$u_q(y_q) - \overline{u_q} = \lambda_q \int_M G(y_q, x) f(x) u_q(x)^{q-1} dv_g(x) \quad (38)$$

for all sequences $y_q \in M$. We write thanks to Lebesgue's dominated convergence theorem, to (29) and to (37) that

$$\begin{aligned} & \int_M G(y_q, x) f(x) u_q(x)^{q-1} \mathbf{1}_{\{u_q \leq \alpha \eta_q\}} dv_g(x) \\ &= \eta_q^{q-1} \int_M G(y_q, x) f(x) dv_g + o\left(\eta_q^{q-1}\right) \\ &= -\frac{2^n \omega_{n-1}}{n \bar{f}} u_q(x_q)^{-1} \int_M G(y_q, x) f(x) dv_g + o\left(u_q(x_q)^{-1}\right) \end{aligned}$$

for all $\alpha > 0$. We can write now that

$$\begin{aligned} \int_M G(y_q, x) f(x) dv_g &= \int_M G(y_q, x) (f(x) - \bar{f}) dv_g \\ &= -\bar{f} \frac{1}{\lambda_2^*} \varphi(y_q) \end{aligned}$$

so that we obtain that

$$\begin{aligned} & \lambda_q \int_M G(y_q, x) f(x) u_q(x)^{q-1} \mathbf{1}_{\{u_q \leq \alpha \eta_q\}} dv_g(x) \\ &= \frac{2^n \omega_{n-1}}{n} u_q(x_q)^{-1} \varphi(y_q) + o\left(u_q(x_q)^{-1}\right). \end{aligned} \quad (39)$$

We assume in the following that $\frac{d_g(x_q, y_q)}{\mu_q} \rightarrow +\infty$ since, otherwise, the estimate of the claim is a straightforward consequence of claim 2.2. We write thanks to claim 2.2 that

$$\begin{aligned} & \int_M G(y_q, x) f(x) u_q(x)^{q-1} \mathbf{1}_{\{u_q > \alpha \eta_q\}} dv_g(x) \\ &= \int_{B_{x_q}(R\mu_q)} G(y_q, x) f(x) u_q(x)^{q-1} dv_g(x) \\ &+ \int_{M \setminus B_{x_q}(R\mu_q)} G(y_q, x) f(x) u_q(x)^{q-1} \mathbf{1}_{\{u_q > \alpha \eta_q\}} dv_g(x). \end{aligned}$$

Then, thanks to the estimate of claim 2.5, to (33) and (37) - note that (37) prevents the second situation from happening in (33), and to standard estimates on the Green

function, we can write that

$$\begin{aligned}
& \int_{M \setminus B_{x_q}(R\mu_q)} G(y_q, x) f(x) u_q(x)^{q-1} \mathbf{1}_{\{u_q > \alpha \eta_q\}} dv_g(x) \\
& \leq C \mu_q^{\frac{n+2}{2}(1-2\varepsilon)} \int_{M \setminus B_{x_q}(R\mu_q)} d_g(y_q, x)^{2-n} d_g(x_q, x)^{(2-n)(1-\varepsilon)(q-1)} dv_g(x) \\
& \quad + C u_q(x_q)^{-1} \int_{M \setminus B_{x_q}(R\mu_q)} d_g(y_q, x)^{2-n} d_g(x_q, x)^{(2-n)\varepsilon} \mathbf{1}_{\{u_q > \alpha \eta_q\}} dv_g(x) \\
& \leq C R^{n-(n+2)(1-\varepsilon)} \mu_q^{\frac{n-2}{2}} d_g(x_q, y_q)^{2-n} + C u_q(x_q)^{-1} \alpha^{-\frac{2-(n+2)\varepsilon}{(n-2)\varepsilon}}
\end{aligned}$$

for some $C > 0$ independent of R , α and q thanks to the fact that $\varepsilon < \frac{2}{n+2}$. In particular, we obtain that

$$\lim_{\alpha \rightarrow +\infty} \lim_{R \rightarrow +\infty} \lim_{q \rightarrow 2^*} \frac{\int_{M \setminus B_{x_q}(R\mu_q)} G(y_q, x) f(x) u_q(x)^{q-1} \mathbf{1}_{\{u_q > \alpha \eta_q\}} dv_g(x)}{u_q(x_q)^{-1} d_g(x_q, y_q)^{2-n}} = 0. \quad (40)$$

It is now easily checked thanks to claim 2.2 that

$$\begin{aligned}
& \lim_{R \rightarrow +\infty} \lim_{q \rightarrow 2^*} \frac{\int_{B_{x_q}(R\mu_q)} G(y_q, x) f(x) u_q(x)^{q-1} dv_g(x)}{u_q(x_q)^{-1} G(y_q, x_q)} \\
& = \frac{\omega_{n-1}}{n} \left(\frac{n(n-2)}{\lambda_{2^*}} \right)^{\frac{n}{2}} f(x_0)^{1-\frac{n}{2}}. \quad (41)
\end{aligned}$$

Coming back to (38) with (39), (40) and (41), we obtain using claim 2.3 that

$$\begin{aligned}
u_q(y_q) - \overline{u_q} &= \left((n-2) \omega_{n-1} \left(\frac{2^n}{\omega_n f(x_0)} \right)^{\frac{n-2}{2}} + o(1) \right) u_q(x_q)^{-1} G(y_q, x_q) \\
&\quad + \frac{2^n \omega_{n-1}}{n} u_q(x_q)^{-1} \varphi(y_q) + o(u_q(x_q)^{-1})
\end{aligned}$$

when $\frac{d_g(x_q, y_q)}{\mu_q} \rightarrow +\infty$. Using claim 2.2, one ends the proof. $\diamond$

We conclude the proof of the theorem. We set

$$R_q = u_q(x_q) (u_q - \overline{u_q})$$

and we write that

$$\Delta_g R_q = \lambda_q u_q(x_q) f u_q^{q-1}.$$

Let K be a compact subset of $M \setminus \{x_0\}$. Thanks to proposition 2.6, we know that

$$R_q \rightarrow 2^n \frac{\omega_{n-1}}{n} \varphi + (n-2) \omega_{n-1} 2^{n-2} f(x_0)^{-\frac{n-2}{n}} G(x_0, \cdot)$$

in $C^0(K)$ as $q \rightarrow 2^*$. Using also (29) and (37), we get that

$$\Delta_g R_q \rightarrow -\frac{2^n \omega_{n-1}}{n} \lambda_{2^*} \frac{f}{f}$$

in $C^1(K)$ as $q \rightarrow 2^*$. Thus, by standard elliptic theory, the above convergence of R_q holds in $C^2(K)$ and we can pass to the limit in the equation satisfied by R_q to

obtain that

$$\begin{aligned} -\frac{2^n \omega_{n-1}}{n} \lambda_{2^*} \frac{f}{\tilde{f}} &= \frac{2^n \omega_{n-1}}{n} \Delta_g \varphi + (n-2) \omega_{n-1} 2^{n-2} f(x_0)^{-\frac{n-2}{n}} \Delta_g G(x_0, \cdot) \\ &= \frac{2^n \omega_{n-1}}{n} \lambda_{2^*} \left(1 - \frac{f}{\tilde{f}}\right) \end{aligned}$$

which is clearly a contradiction. This proves the theorem.

Remarks : the solution we obtain is a minimizing solution in the sense we have that

$$\int_M |\nabla u|_g^2 dv_g = \lambda_{2^*}(f)$$

and $u \in \mathcal{H}_{2^*}$ where $\lambda_{2^*}(f)$ and $\mathcal{H}_{2^*}(f)$ are as in claim 2.1. We add the dependence in f with respect to the notations of claim 2.1. Note also that as a consequence of this result, we have that

$$\lambda_{2^*}(f) < K(n, 2)^{-2} \left(\max_M f \right)^{-\frac{2}{2^*}} \quad (42)$$

for all smooth functions f satisfying (5) and (6), a result which is not obvious a priori. Indeed, imagine that, on the contrary, there exists some $f \in C^\infty(M)$ satisfying (5) and (6) and such that

$$\lambda_{2^*}(f) = K(n, 2)^{-2} \left(\max_M f \right)^{-\frac{2}{2^*}}. \quad (43)$$

Let us look at the function $\tilde{f} = -\max_M f + 2f$ which satisfies (5) and (6) and which satisfies moreover that $\max_M \tilde{f} = \max_M f$. By test functions computations we know that

$$\lambda_{2^*}(\tilde{f}) \leq K(n, 2)^{-2} \left(\max_M f \right)^{-\frac{2}{2^*}}.$$

Moreover, thanks to what we proved above, there exists $u \in C^\infty(M)$, $u > 0$, which satisfies that

$$\int_M \tilde{f} u^{2^*} dv_g = 1$$

and that

$$\int_M |\nabla u|_g^2 dv_g = \lambda_{2^*}(\tilde{f}) \leq K(n, 2)^{-2} \left(\max_M f \right)^{-\frac{2}{2^*}}.$$

Since $f \geq \tilde{f}$ and $f \not\equiv \tilde{f}$, we have that

$$\int_M f u^{2^*} dv_g > 1$$

and we can write thanks to the definition of $\lambda_{2^*}(f)$ that

$$\lambda_{2^*}(f) \left(\int_M f u^{2^*} dv_g \right)^{\frac{2}{2^*}} \leq \int_M |\nabla u|_g^2 dv_g \leq K(n, 2)^{-2} \left(\max_M f \right)^{-\frac{2}{2^*}}.$$

This is clearly in contradiction with (43). Thus the above claim (42) is proved.

Acknowledgements : I would like to thank Emmanuel Hebey and Frédéric Robert for their interest in this work and their fruitful comments about this paper.

REFERENCES

1. H. Berestycki, L. Nirenberg, and S. Varadhan, *The principal eigenvalue and maximum principle for second order elliptic operators in general domains*, Comm. Pure Appl. Math. **47** (1994), 47–92.
2. H. Brézis and E. Lieb, *A relation between pointwise convergence of functions and convergence of functionals*, Proc. Amer. Math. Soc. **88**, **3** (1983), 486–490.
3. L.A. Caffarelli, B. Gidas, and J. Spruck, *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth*, Comm. Pure Appl. Math. **42** (1989), 271–297.
4. O. Druet, *The best constants problem in Sobolev inequalities*, Math. Ann. **314** (1999), 327–346.
5. O. Druet, E. Hebey, and F. Robert, *Blow-up theory for elliptic PDEs in Riemannian geometry*, Mathematical Notes, vol. 45, Princeton University Press, 2004.
6. J.F. Escobar and R. Schoen, *Conformal metrics with prescribed scalar curvature*, Inventiones Mathematicae **86** (1986), 243–254.
7. D. Gilbarg and N. Trudinger, *Elliptic partial differential equations of second order*, Springer-Verlag, Berlin Heidelberg New York, 1977.
8. E. Hebey and M. Vaugon, *The best constant problem in the Sobolev embedding theorem for complete Riemannian manifolds*, Duke Math. J. **79** (1995), 235–279.
9. ———, *Meilleures constantes dans le théorème d’inclusion de Sobolev*, Ann. Inst. H. Poincaré. Anal. Non Linéaire **13** (1996), 57–93.
10. Y.T. Jung, *On the elliptic equation $\frac{4(n-1)}{n-2}\Delta u + K(x)u^{\frac{n+2}{n-2}} = 0$ and the conformal deformation of Riemannian metrics*, Indiana Univ. Math. J. **43** (1994), 737–746.
11. J.L. Kazdan, *Prescribing the curvature of a Riemannian manifold*, CBMS, vol. 57, A.M.S., Providence, R.I., 1984.
12. J.L. Kazdan and F.W. Warner, *Curvature function for compact 2-manifolds*, Ann. of Math. **99** (1974), 14–74.
13. ———, *Scalar curvature and conformal deformation of Riemannian structure*, Journal of Differential Geometry **10** (1975), 113–134.
14. N.S. Trudinger, *Remarks concerning the conformal deformation of Riemannian structures on compact manifolds*, Ann. Scuola Norm. Sup. Pisa **22** (1968), 265–274.
15. H. Yamabe, *On a deformation of Riemannian structures on compact manifolds*, Osaka Math. J. **12** (1960), 21–37.

OLIVIER DRUET, UMPA - ENS LYON, 46, ALLÉE D’ITALIE, F-69364 LYON CEDEX 7, FRANCE
E-mail address: odruet@umpa.ens-lyon.fr